

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau

PCT

(43) International Publication Date
11 July 2002 (11.07.2002)(10) International Publication Number
WO 02/053133 A1

PCT

(31) International Patent Classification: A61K 9/16,
C07D 417/87, A61P 25/24, A61K 9/14 // 31/34

(21) International Application Number: PCT/DK02/00003

(22) International Filing Date: 3 January 2002 (03.01.2002)

(23) Filing Language: English

(24) Publication Language: English

(30) Priority Data:
PA2001 00016 3 January 2001 (05.01.2001) DK(71) Applicant (for all designated States except US): H.
LUNDBECK A/S (DK/DK); (Købmøllevej 9, DK-2500
Valby-Copenhagen (DK).

(72) Inventors; and

(73) Inventors/Applicants (for US only): LILJEGRÉN, Ken
(DK/DK); Mellemsten 30, Jernstrup, DK-1500 Varde
(DK). HOLM, Per (DK/DK); Ørstedals Parkvej 54,
DK-2720 Vanløse (DK).(81) Designated States (national): AE, AG, AL, AM, AT, AT
(utility model), AU, A7, BA, BB, BG, BR, BY, BZ, CA,
CH, CN, CO, CR, CU, CZ, CZ (utility model), DE, DE
(utility model), DK, DK (utility model), DM, DZ, EC, EE,
EE (utility model), ES, FI, FI (utility model), GB, GD, GR,
GH, GM, HR, HU, IL, IL, IN, IS, JP, KE, KG, KP, KR,
KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK,
MN, MW, MX, MY, NZ, NZ, OM, PH, PL, PT, RO, RU,
SD, SE, SG, SI, SK, SK (utility model), SL, TJ, TM, TN,
TR, TT, TZ, UA, UA, US, UZ, VN, YU, ZA, ZM, ZW.(84) Designated States (regional): ARIPO (BREM) (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR,
GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent
(BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR,
NE, SN, TD, TG).

Published:

— with international search report

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

WO 02/053133 A1

(54) Title: PHARMACEUTICAL COMPOSITION CONTAINING CITALOPRAM

(57) Abstract: A solid unit dosage form comprising citalopram which is prepared by a process comprising a step wherein citalopram
base or a pharmaceutically acceptable salt and optionally pharmacologically acceptable excipients is roller compacted.

WO 02/053133


PCT/DK02/00003

Pharmaceutical composition containing Citalopram

The present invention relates to a novel pharmaceutical composition containing citalopram, 1-[3-(dimethylamino)propyl]-1-(4-fluorophenyl)-1,3-dihydro-5-isobenzofurancarbonitrile.

Background of the Invention.

Citalopram is a well-known antidepressant drug that has the following structure:



It is a selective, centrally active serotonin (5-hydroxytryptamine; 5-HT) reuptake inhibitor, accordingly having antidepressant activities.

Citalopram was first disclosed in DE 2,657,013, corresponding to US 4,136,193. This patent publication describes the preparation of citalopram by one method and outlines a further method which may be used for preparing citalopram. The citalopram prepared was isolated in crystalline form as the oxalate, the hydrobromide and the hydrochloride salt, respectively. Furthermore, the citalopram base was obtained as an oil (B.P. 175 C/0.03 mmHg). The publication also outlines the manufacture of tablets containing salts of citalopram. Citalopram is marketed as the hydrobromide and the hydrochloride, respectively.

Manufacture of crystalline citalopram base is disclosed in co-pending DK 2000 00402. This patent publication describes the preparation of crystalline citalopram base and the use of crystalline citalopram base as an intermediate in the purification of crude citalopram hydrobromide into pure citalopram hydrobromide. The publication also outlines the manufacture of tablets containing citalopram base.

CONFIRMATION COPY

WO 02/033133

PCT/DK02/00003

2

Citalopram is marketed in a number of countries as a tablet prepared by compression of wet-granulated citalopram hydrobromide, lactose and other excipients.

5 It is well-recognised that preparation of tablets with a reproducible composition requires that all the dry ingredients have good flow properties. In cases, where the active ingredient has good flow properties, tablets can be prepared by direct compression of the ingredients. However, in many cases, where the particle size of the active substance is small, the active substance is cohesive or has poor flow properties.

10 Further, active substances with a small particle size mixed with excipients having a larger particle size will typically segregate or de-mix during the tableting process.

15 The problems of small particle size, poor flowability and segregation are conventionally solved by enlarging the particle size of the active substance, usually by granulation of the active ingredient either alone or in combination with a filler and/or other conventional tablet ingredients.

20 One such granulation method is the "wet" granulation process. Using this method, the dry solids (active ingredients, filler, binder etc.) are blended and moistened with water or another wetting agent (e.g. an alcohol) and agglomerates or granules are built up of the moistened solids. Wet massing is continued until a desired homogenous particle size has been achieved whereupon the granulated product is dried.

25 An alternative to the "wet" granulation method is the "melt" granulation, which is also known as the "thermal plastic" granulation process, where a low melting solid is used as the granulation agent. Initially, the dry solids are blended and heated until the binder melts. As the binder is liquefied and spreads over the surface of the particles, the particles will adhere to each other and form granules. The binder solidifies upon cooling forming a dry granular product.

30 Wet granulation as well as melt granulation are energy intensive unit operations requiring complicated and expensive equipment as well as technical skill.

WO 02/053133

PCT/DK02/00003

3

The process used for the preparation of citalopram hydrobromide results in a product with a very small particle size around 2-20 μm that, as many other particulate products with a small particle size, has very poor flow properties. Thus, in order to achieve appropriate dosing of the citalopram during tableting, it is considered
5 necessary to make a granulate of citalopram with larger particle size and improved flow properties.

The citalopram tablet that is marketed is a tablet made from fluid-bed dried, wet-
10 granulated citalopram hydrobromide with various excipients.

A third size enlargement method is roller compaction where the size enlargement is done by mechanical means. Using this method, the dry solids are compressed between two rollers resulting in a sheet which subsequently is broken down into a granulate by mechanical means such as a rotating mill and oscillating screens.
15

The integration of the granulation into one apparatus in roller compaction results in that the process is difficult to control and tends to give very broad or even bimodal particle size distributions. Broad or bimodal particle size distributions will often have adverse effects, such as poor flow characteristics, segregation, de-mixing and the like,
20 hampering the later stages of the formulation of a pharmaceutical acceptable solid unit dosage form with constant composition.

In view of the fact that roller compaction requires fewer process steps, is much less time consuming and cheaper than the processes involving wet or melt granulation,
25 there is a desire for a process for roller compaction of citalopram hydrobromide.

The obstacles that hitherto have hindered roller compaction of citalopram tablets have now been circumvented.

30 It has, surprisingly, been found that a granulate prepared by roller compaction of essentially undiluted citalopram and having a median particle size comparable to the median particle size of the filler is useful for the manufacture of compressed tablets despite the broad or bimodal particle size distribution of the granulate.

WO 02/053133

PCT/DK02/00003

4

Likewise surprising, it has been found that a granulate prepared by roller compaction of citalopram mixed with all excipients for the finished formulation except for a small amount of glidant is useful for the manufacture of compressed tablets despite the broad or bimodal particle size distribution of the granulate.

- 1 Accurate dosing in capsules may also be with such roller compacted granulates.

Objects of the Invention

- 10 It is the object of the present invention to provide a novel pharmaceutical unit dosage form containing roller compacted citalopram.

A second object of the invention is to provide a capsule containing citalopram.

- 15 A third object of the invention is to provide a roller compacted granulate comprising citalopram.

A fourth object of the invention is to provide a process for roller compaction of citalopram.

20 Summary of the Invention

The invention then, *inter alia*, comprises the following alone or in combination:

- 25 A solid unit dosage form comprising citalopram prepared by roller compaction of citalopram base or a pharmaceutically acceptable salt thereof, where pharmaceutically acceptable excipients optionally may be mixed with the active ingredient before granulation, and optionally the roller compacted granulate may be mixed with extragranular pharmaceutically acceptable excipients, whereupon said granulate or mixture with extragranular excipients is compressed into a tablet or filled in a hard
30 gelatine capsule.

A granulate comprising citalopram base or a pharmaceutically acceptable salt thereof, where said granulate is formed by roller compaction of a powder comprising

WO 02/053133

PCT/DK02/00003

5

citalopram base or a pharmaceutically acceptable salt thereof and optionally pharmaceutically acceptable excipients.

5 A method for manufacture of a granulate comprising citalopram base or a pharmaceutically acceptable salt thereof, where said method comprises roller compaction of a powder comprising citalopram base or a pharmaceutically acceptable salt thereof and optionally pharmaceutically acceptable excipients.

10 Citalopram can be compacted alone or optionally mixed with a small amount of glidant, such as magnesium stearate, to minimize adhesion to surfaces in the compaction equipment. Afterwards, the granulate is mixed with extragranular excipients in order to form a mixture, which can be compressed into a tablet or filled in a hard gelatine capsule.

15 At the other end of the scale, citalopram may be mixed with all excipients prior to compaction, or, optionally, all ingredients but a small amount of glidant, which is added after compaction. Thus, the granulate, optionally admixed with glidant, is ready for tableting or filling in a hard gelatine capsule. All ingredients are "locked" in the granule and cannot detach.

20 The roller compaction of citalopram and optional pharmaceutically acceptable excipients into a granulate, which can be used in formulation of pharmaceutical acceptable solid unit dosage forms has the great advantage, that wet or melt granulation, which requires a time-consuming heating or drying step, is avoided.

25 As used herein, "particle size distribution" means the distribution of equivalent spherical diameters as determined by laser diffraction in a Sympatec Helos equipment. The particle size distributions for fillers and uncompacted citalopram are determined at 1 bar dispersive pressure, whereas the particle size distributions for compacted granulates are determined at 0.2 bar dispersive pressure in order to avoid deaggregation of the granules leading to erroneous results. "Median particle size",
30 correspondingly, means the median of said particle size distribution.

WO 02/053133

PCT/DK02/00003

6

Thus in one embodiment of the invention, the present invention relates to a tablet prepared by compression of a mixture of roller compacted citalopram base or a pharmaceutically acceptable salt thereof and pharmaceutically acceptable excipients.

- 5 In another embodiment, the present invention relates to a capsule prepared by filling a mixture of roller compacted citalopram base or a pharmaceutically acceptable salt thereof and pharmaceutically acceptable excipients in a hard gelatine capsule.

- 10 Flow, segregation and demixing properties and, hence, the suitability of the granulates for compression into tablets or filling in hard gelatine capsules depend, besides the median particle size, on the particle size distribution.

Preferably, the solid unit dosage forms according to the invention do not contain a binder.

15

- The solid unit dosage form according to the invention may contain 2-60% w/w active ingredient calculated as citalopram base, preferably 10-40% w/w active ingredient calculated as citalopram base, and more preferred 15-25% w/w active ingredient calculated as citalopram base. Suitably, the solid unit dosage form of the invention
20 contains 20% w/w active ingredient calculated as citalopram base.

- In one preferred embodiment of the invention, the present invention relates to a solid unit dosage form wherein the active ingredient is citalopram hydrobromide or citalopram hydrochloride. Preferably the active ingredient contained in the solid unit
25 dosage form of the invention is citalopram hydrobromide.

In another preferred embodiment of the invention, the present invention relates to a solid unit dosage form wherein the active ingredient is citalopram base.

- 30 The solid unit dosage form according to the invention may contain a filler selected from lactose, or other sugars e.g. sorbitol, mannitol, dextrose and sucrose, calcium phosphates (dibasic, tribasic, hydrous and anhydrous), starch, modified starches,

WO 02/053133

PCT/DK02/00003

7

microcrystalline cellulose, calcium sulphate and/or calcium carbonate. In a preferred embodiment, the solid unit dosage form of the invention does not contain lactose.

Suitably the filler is a microcrystalline cellulose such as ProSolv SMCC90 manufactured by Penwest Pharmaceuticals or Avicel PH 200 or Avicel PH 101
5 manufactured by FMC Corporation.

Besides the active ingredient and filler, the solid pharmaceutical unit dosage forms may include various other conventional excipients such as disintegrants, and optionally minor amounts of lubricants, colorants and sweeteners.

10

Lubricants used according to the invention may suitably be one or more of the following metallic stearates (magnesium, calcium, sodium), stearic acid, wax, hydrogenated vegetable oil, talc and colloidal silica.

15 Suitably the lubricant is magnesium stearate or calcium stearate

Disintegrants include sodium starch glycolate, croscarmellose, crospovidone, low substituted hydroxypropylcellulose, modified cornstarch, pregelatinized starch and natural starch.

20

The granulate comprising the active ingredient after compaction has preferably a median particle size of at least 40 μm , more preferred in the range of 40 - 250 μm , even more preferred in the range of 45 - 200 μm and most preferred in the range of 50 - 180 μm .

25

The active ingredient is prior to compaction in the form of a powder, which preferably has a median particle size below 20 μm and more preferred below 15 μm .

The solid, pharmaceutical unit dosage form of the invention may be prepared by
30 conventional methods using a tablet press with forced feed capability.

The filled, hard gelatine capsule of the invention may be prepared by conventional methods using a capsule filler suitable for powder filling.

WO 02/053133

PCT/DK02/00003

8

The crystals of a pharmaceutically acceptable salt of citalopram used in one embodiment of the invention may be produced according to methods described in US 4,136,193.

- 5 The crystals of citalopram base used in one embodiment of the invention may be produced according to methods described in co-pending DK 2000 00402.

In the following, the invention is illustrated by way of examples. However, the examples are merely intended to illustrate the invention and should not be construed as limiting.

10

Example 1

Compaction of citalopram hydrobromide

- 15 Citalopram hydrobromide (8000 g) was mixed with Mg-stearate (80 g) by conventional mixing. The mixture was compacted on an Alexanderwerk WP120 x 40 V roller compactor.

The parameters for the compaction were set as follows:

- 20 Roller speed: 8 rpm
Roller pressure: 6.5 kN/cm² (70 bar)
Auger speed: 35 rpm
Product flow: 14 kg/h
Screens: 2.0 mm and 0.8 mm
25 Vacuum: On

The resulting granulate constitutes the intragranular phase in subsequent tableting in example 3. The granulate had the following properties:

- 30 Bulk density: 0.40 g/mL
Tapped density (1250 raps): 0.52 g/mL
Flowability through 15 mm orifice: 5.3 g/s

WO 02/033133

PCT/DK02/00003

9

The particle size distributions for the citalopram hydrobromide used as feed as well as the resulting granulate are listed in table 1.

5 Example 2

Compaction of all ingredients, except magnesium stearate

10 Citalopram hydrobromide (3740 g), Kollidon VA64 (748 g) as binder and Avicel PH 101 (14209 g) as filler was mixed by conventional mixing. The mixture was compacted on an Alexanderwerk WP 200 x 75 V roller compactor.

The parameters for the compaction were set as follows:

Roller speed: 6 rpm

15 Roller pressure: 7,8 kN/cm² (90 bar)

Auger speed: 45 rpm

Product flow: 65 kg/h

Screens: 2.0 mm and 0.6 mm (100 and 70 rpm respectively)

Vacuum: On

20

The resulting granulate constitutes the intragranular phase in subsequent tableting in example 4. The granulate had the following properties:

Bulk density: 0.55 g/mL

25 Tapped density (1250 taps) 0.75 g/mL

The particle size distributions for the feed materials as well as the resulting granulate are listed in table 1.

30

WO 02/053133

PCT/DK02/00003

10

Table 1: Particle size distribution (Sympatec Helos) for citalopram hydrobromide crystals (feed to compaction); compacted material, examples 1 and 2; and excipients, Kollidon VA 64, Avicel PH 101 and ProSolv SCMC90

Quantile (%)	Citalopram HBr (μm)	Example 1 (μm)	Example 2 (μm)	Kollidon VA 64 (μm)	Avicel PH 101 (μm)	ProSolv SCMC90 (μm)
95	97.0	737	712	—	178	280
90	72.3	652	598	148	149	232
50	14.0	169	71.4	63.3	68.5	114
10	1.2	6.3	12.0	18.5	23.4	32.1

Example 3

Tabletting of compacted citalopram hydrobromide mixed with extragranular excipients.

Compacted material (5800 g) from example 1 was mixed with silicified microcrystalline cellulose (ProSolv SMCC90) (22765 g) as filler in a Bohle PTM 200 (100 L) mixer for 3 minutes at 7 rpm. Magnesium stearate (144 g) was added as extra glidant and mixing continued for 30 seconds.

25 kg of the above mixture was tableted on a Fette P 1200 IC tablet press at speeds of 50,000 to 125,000 tablets/hour. The granulate was fed by means of a forced feeder. Tablet core weight was 125 mg corresponding to a tablet strength of 20 mg citalopram base-equivalent.

During tabletting, samples were withdrawn at every 500 g granulate corresponding to every 4000 tablets. Tabletting ended after manufacture of 184,000 tablets.

WO 02/059133

PCT/DK02/00003

11

Two tablets from each sample were assayed by a validated method using UV-absorption in an aqueous solution, thus analysing in total 92 tablets. The relative standard deviation in citalopram content was 4.4%.

5

Example 4

Tabletting of compacted mixture of citalopram hydrobromide, Kollidon VA64 and Avicel PH 101 with extragranular magnesium stearate.

10

Granulate from example 2 was mixed with Mg-stearate as glidant.

Mixing was performed in a Bohle PTM 200 (100 L) mixer for 30 seconds at 7 rpm.

Intragranular phase	%-intragran.	qty (g)	% pr.tab.	mg pr.tab.
Citalopram HBr	20.0 %	3740	19.9 %	25.0
Kollidon VA64	4.0 %	748	4.0 %	5.0
Avicel PH101	76.0 %	14209	75.6 %	95.0
Extragranular phase				
Mg-stearate	0.3%	90	0.3 %	0.6

Table 2: Composition of tablets

15

18 kg of the above mixture was tableted on a Fette P 1200 IC tablet press at speeds of 50,000 to 125,000 tablets/hour. The granulate was fed by means of a forced feeder. Tablet core weight was 125 mg corresponding to a tablet strength of 20 mg citalopram base-equivalent.

20

During tabletting, samples were withdrawn at every 500 g granulate corresponding to every 4000 tablets. Tabletting ended after manufacture of 124,000 tablets.

Two tablets from each sample were assayed by a validated method using UV-absorption in an aqueous solution, thus analysing in total 92 tablets. The relative standard deviation of content of citalopram base equivalent content was 1.2%.

25

WO 02/053133

PCT/DK02/00003

12

Claims

1. A solid unit dosage form comprising citalopram, characterised in that it is prepared by a process comprising a step wherein citalopram base or a pharmaceutically acceptable salt, and optionally pharmaceutically acceptable excipients is roller compacted.
2. The solid unit dosage form according to claim 1, characterised in that the active ingredient is
- a) essentially undiluted at the roller compacting step; or
- b) mixed with essentially all the excipients at the roller compacting step.
3. The solid unit dosage form according to claims 1-3, characterised in that it contains 2-60% w/w active ingredient calculated as citalopram base, preferably 10-40% w/w active ingredient calculated as citalopram base and more preferred 15-25% w/w active ingredient calculated as citalopram base.
4. The solid unit dosage form according to claim 1-3, characterised in that the granulate comprising the active ingredient after compaction has a median particle size of at least 40 μm , preferably in the range of 40 - 250 μm , more preferred in the range of 45 - 200 μm and most preferred in the range of 50 - 180 μm .
5. The solid unit dosage form of claims 1-4, characterised in that the active ingredient is citalopram hydrobromide or citalopram hydrochloride.

1

INTERNATIONAL SEARCH REPORT

International application No.
PCT/DK 02/00009

A. CLASSIFICATION OF SUBJECT MATTER

IPC7: A61K 9/16, C07D 307/87, A61P 25/24, A61K 9/14 // A61K 31/34
According to International Patent Classification (IPC) or to both national classification and IPC:

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC7: A61K, C07D, A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

SE, DK, FI, NO classes as above

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P, X	EP 1152000 A1 (SUMIKA FINE CHEMICALS CO. LTD.), 7 November 2001 (07.11.01), claims 1-19 --	1-5
P, X	WO 0180619 A2 (H. LUNDBECK A/S), 1 November 2001 (01.11.01), claims 1-33 --	1-5
X	Churchill Livingstone, "Pharmaceutics: The Science of dosage form design", Aulton Michael, 1998, pages 627-628 --	1-5
A	GB 1358915 A (MERCK & CO. INC.), 3 July 1974 (03.07.74), claims 1-13 --	1-5

☒ Further documents are listed in the continuation of Box C.☒ See patent family annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"C" earlier application or patent not published on or after the international filing date

"L" document which may throw doubt on priority claim(s) or which is cited to establish the publication date of another citation or other special reasons (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"I" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"R" document member of the same patent family

Date of the actual completion of the international search

2 April 2002

Date of mailing of the international search report

11-04-2002

Name and mailing address of the ISA/
Swedish Patent Office
Box 3055, S-102 42 STOCKHOLM
Facsimile No. +46 8 666 02 86

Authorized officer

FERNANDO FARIETA/BS
Telephone No. +46 8 782 25 00

Form PCT/ISA/210 (second sheet) (July 1998)

2

INTERNATIONAL SEARCH REPORT

International application No.
PCT/DK 02/00003

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,A	US 2001/0031784 A1 (PETERSEN ET AL), 18 October 2001 (18.10.01), claims 1-12 --	1-5
A	US 4136193 A (BOGESD ET AL), 23 January 1979 (23.01.79), claims 1-18 -- -----	1-5

INTERNATIONAL SEARCH REPORT

Information on patent family members

28/01/02

International application No.

PCT/DK 02/00003

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 1152000 A1	07/11/01	AU 3508501 A	08/11/01
		US 2001049450 A	06/12/01
WO 0180619 A2	01/11/01	DE 20113195 U	15/11/01
		DK 200100203 U	10/11/01
		FI 5176 U	21/12/01
		FR 2812811 A	15/02/02
		GB 0118579 D	00/00/00
		HU 0103071 D	00/00/00
		NO 20013891 D	00/00/00
GB 1358915 A	03/07/74	NONE	
US 2001/0031784 A1	18/10/01	NONE	
US 4136193 A	23/01/79	AT 959488 B	10/11/80
		AT 360001 B	10/12/80
		AT 360002 B	10/12/80
		AT 571979 A	15/05/80
		AT 572079 A	15/05/80
		AT 947276 A	15/04/80
		AU 509445 B	15/05/80
		AU 2107377 A	13/07/79
		BE 850401 A	14/07/77
		CA 1094087 A	20/01/81
		CH 626886 A	15/12/81
		CH 632258 A	30/09/82
		CH 632259 A	30/09/82
		DE 2657013 A,C	28/07/77
		UK 13177 A	15/07/77
		DK 143275 B,C	03/08/81
		ES 454980 A	01/04/78
		FI 63754 B,C	29/04/83
		FI 770073 A	15/07/77
		FR 2938271 A,B	12/08/77
		GB 1526331 A	27/09/78
		IE 44055 B	29/07/81
		JP 1368581 C	11/03/87
		JP 52105162 A	03/09/77
		JP 61035986 B	15/08/86
		NL 192451 B,C	01/04/97
		NL 7700244 A	18/07/77
		NO 147243 B,C	22/11/82
		NO 770109 A	15/07/77
		NZ 183001 A	02/06/78
		SE 429551 B,C	12/09/83
		SE 7614201 A	15/07/77
		ZA 7700057 A	30/11/77

*** RX REPORT ***

RECEPTION OK

TX/RX NO	7942
RECIPIENT ADDRESS	212 753 6237
DESTINATION ID	
ST. TIME	01/28 13:21
TIME USE	04'16
PGS.	24
RESULT	OK